

New findings of *Neurospora* in Europe and comparisons of diversity in temperate climates on continental scales

David J. Jacobson¹

Department of Biological Sciences, Stanford University, Stanford, California 94305-5020 and Department of Plant and Microbial Biology, University of California, Berkeley, California 94720-3102

Jeremy R. Dettman²

Rachel I. Adams³

Department of Plant and Microbial Biology, University of California, Berkeley, California 94720-3102

Cornelia Boesl

Shahana Sultana⁴

Till Roenneberg

Martha Merrow⁵

Institute for Medical Psychology, Centre for Chronobiology, Goethestraße 31, Ludwig Maximilians University München, 80336 München, Germany

Margarida Duarte

Isabel Marques

Alexandra Ushakova

Patrícia Carneiro

Arnaldo Videira

Instituto de Biologia Molecular e Celular (IBMC), and Instituto de Ciências Biomédicas Abel Salazar, (ICBAS) Universidade do Porto, Portugal

Laura Navarro-Sampedro

María Olmedo

Luis M. Corrochano

Departamento de Genética, Facultad de Biología, Universidad de Sevilla, Avenida Reina Mercedes 6, Apartado 1095, E-41080 Sevilla, Spain

John W. Taylor

Department of Plant and Microbial Biology, University of California, Berkeley, California 94720-3102

Abstract: The life cycles of the conidiating species of *Neurospora* are adapted to respond to fire, which is reflected in their natural history. *Neurospora* is found commonly on burned vegetation from the tropic and

subtropical regions around the world and through the temperate regions of western North America. In temperate Europe it was unknown whether *Neurospora* would be as common as it is in North America because it has been reported only occasionally. In 2003 and 2004 a multinational effort surveyed wildfire sites in southern Europe. *Neurospora* was found commonly from southern Portugal and Spain (37°N) to Switzerland (46°N). Species collected included *N. crassa*, *N. discreta*, *N. sitophila* and *N. tetrasperma*. The species distribution and spatial dynamics of *Neurospora* populations showed both similarities and differences when compared between temperate Europe and western North America, both regions of similar latitude, climate and vegetation. For example the predominant species in western North America, *N. discreta* phylogenetic species 4B, is common but not predominant in Europe, whereas species rare in western North America, *N. crassa* NcB and *N. sitophila*, are much more common in Europe. The meiotic drive element Spore killer was also common in European populations of *N. sitophila* and at a higher proportion than anywhere else in the world. The methods by which organisms spread and adapt to new environments are fundamental ecosystem properties, yet they are little understood. The differences in regional diversity, reported here, can form the basis of testable hypotheses. Questions of phylogeography and adaptations can be addressed specifically by studying *Neurospora* in nature.

Key words: ecology, fire, meiotic drive, natural history, phylogenetic species, Spore killer

INTRODUCTION

The conidiating species of the ascomycete fungus *Neurospora*, as a group, have been considered to be primarily tropical or subtropical with a complete longitudinal distribution (Turner and Perkins 1988, Turner et al 2001). These particular *Neurospora* species are well adapted to grow and sporulate on the surface of fire-scorched vegetation. Recent field surveys, however, have found that *Neurospora* commonly occupies an entirely different ecological niche, in dry and/or cold habitats. Within this new geographic range, western North America from New Mexico (34°N) to Alaska (64°N) (Jacobson et al 2004), *Neurospora* was found under the bark of fire-damaged trees. This discovery has raised questions

Accepted for publication 14 Jul 2006.

¹ Corresponding author. E-mail: djacob@stanford.edu

² Present address: Department of Botany, University of Toronto, Mississauga, Ontario, L5L 1C6 Canada.

³ Present address: Department of Biological Sciences, Stanford University, Stanford, California 94305-5020.

⁴ Present address: Internal Medicine I, Klinikum Grosshadern Marchioninstr 15, Ludwig Maximilians University München, 81377 München, Germany.

⁵ Present address: Biological Centre, Chronobiology, University of Groningen, Postbus 14, 9750AA Haren, The Netherlands.

about the occurrence of *Neurospora* in other temperate regions.

The purpose of this study was to determine whether *Neurospora* is common in temperate regions of Europe. We hypothesized that the niche under the bark of burned vegetation had been overlooked in Europe as it was in North America. In autumn 2003 a multinational effort searched for *Neurospora* in fire sites across southern Europe after a summer of unusually devastating wildfires. Additional collections were made in 2004.

Most published accounts of *Neurospora* in temperate regions were anecdotal (see Jacobson et al 2004). In Europe *Neurospora* most often has been associated with bakeries, (Legan 1993, Perkins 1991, Perkins and Turner 1988, Yassin and Wheals 1992). High temperatures and the presence of easily colonized substrates that usually are associated with bakeries may allow *Neurospora* to grow in locations that traditionally were considered outside the geographic distribution of this fungus. However observations of *Neurospora* in nature have been sporadic in Europe with no systematic surveys or descriptions of populations on the scale of studies in temperate North America.

Individuals collected in Europe were identified with both biological and phylogenetic species recognition methods that have been developed for the outbreeding species of *Neurospora*. Phylogenetic species recognition also provided a preliminary indication of genetic diversity within species. The comparison of the isolates collected in this study with those from North America and throughout the world highlights differences in the ecology of *Neurospora* and the diversity of *Neurospora* populations in temperate climates on different continents.

MATERIALS AND METHODS

Collection, culturing and species identification.—An international consortium was formed to survey for *Neurospora* in Europe in summer 2003 during which an unusually intense heat wave led to devastating wildfires. Fire progression was followed and fire maps obtained from the Global Fire Monitoring Center website, <http://www.fire.uni-freiburg.de/current/globalfire.htm>, and links therein. Satellite photos were obtained when possible from the Moderate Resolution Imaging Spectroradiometer (MODIS) Rapid Response system, near-real-time production website, <http://rapidfire.sci.gsfc.nasa.gov/production>. These maps and photographs were used to locate easily accessible and widely distributed sites across southern Europe. Initial surveys were made in Portugal in early Sep 2003. Systematic field work was conducted in late Sep and early Oct 2003 at sites in Switzerland, northern Italy, southern France

and northern Spain. Additional collections from Sapiãos, Portugal, and Seville, Spain, were made respectively in Sep and Oct 2004.

Methods of handling isolates, including collecting, initial culturing, subculturing of single conidia and storage, were exactly as described in Jacobson et al (2004). A field sample of conidia was collected from a sporulating colony onto sterile filter paper, which then was placed in a sterile envelope. One colony per plant was sampled for up to 45 isolates per site. In addition, where possible, two to seven isolates from the same plant were collected from one or two plants per site. Representative isolates of each species found at each site (both mating types when possible), and strains (TABLE I see below) have been deposited in the Fungal Genetic Stock Center (FGSC), Kansas city, Missouri 64110 (<http://www.fgsc.net>) under accession numbers 10010–10059.

Recent taxonomic work (Cai et al 2006, García et al 2004) has not changed the status of the conidiating species of *Neurospora*. Therefore biological species recognition was used to identify isolates to species with a three-step process following methods outlined in Perkins and Turner (1988): (i) Assessment of heterothallism: A single conidium subculture from each isolate was allowed to grow on Vogel's minimal medium N (Davis 2000) at 25 C for 7–10 d to test for self-fertility. Perithecia from each self-fertile isolate were dissected to determine the number of ascospores per ascus. All isolates with four ascospores per ascus were concluded to be *N. tetrasperma*. (ii) Mating-type (*mat*) determination: Each self-sterile (heterothallic) isolate was crossed to both *mat A* and *mat a* species tester strains of *N. crassa* that contain the fluffy mutation (FGSC strains 6682 and 6683, respectively). Conidia of unknown isolates (males) were used to fertilize protoperithecia of the tester strains (females) growing on Petri dishes of Westergaard's synthetic crossing medium (Davis 2000). Fertilization was successful when conidia were applied to a small region of the tester colony, so that a single female tester on a 9 cm diam plate could be fertilized with up to 30 different isolates. A darkening and swelling of protoperithecia indicated a mating reaction after 2–4 d incubation at 25 C. A positive mating reaction on one female tester was obtained for each isolate, thus revealing mating type. (iii) Mating with species testers: When crossed to the *N. crassa* tester any isolate that produced >50% black ascospores after 7–10 d postfertilization was classified as *N. crassa* (Perkins and Turner 1988). Isolates that produced only hyaline, unviable ascospores or no spores at all were judged not to be *N. crassa*. This response also confirmed that none of these isolates were *N. intermedia*, which routinely produces 5–10% black ascospores with the *N. crassa* tester strains (Perkins and Turner 1988). Each isolate was crossed, again as a male, to plates of tester strain females of the appropriate mating type for both *N. sitophila* (FGSC 5940 *mat A* or 5941 *mat a*) and *N. discreta* (FGSC 3228 *mat A* or 4378 *mat a*). Fertility to these testers was mutually exclusive. Production of black ascospores was limited to crosses with one and only one of the species testers; no isolate made black ascospores with more than one tester. In addition no isolate was infertile with all *Neurospora* species testers.

TABLE I. European isolates of *Neurospora* used in phylogenetic analyses

Species, clade and isolate numbers			
FGSC number	D number ^a	Mating type	Country, site
<i>N. crassa</i> NcB			
10049		<i>mat A</i>	Spain, Platja d'Aro
10050		<i>mat A</i>	Spain, Platja d'Aro
10033		<i>mat a</i>	Spain, Macanet de la Selva
10043		<i>mat a</i>	Spain, Seros
10044		<i>mat A</i>	Spain, Seros
10045		<i>mat a</i>	Spain, Seros
10046		<i>mat A</i>	Spain, Seros
10017		<i>mat A</i>	Portugal, Troviscal Sertã
10018		<i>mat A</i>	Portugal, Penedo Furado
10020		<i>mat A</i>	Portugal, Tapada de Mafra
10021		<i>mat a</i>	Portugal, Tapada de Mafra
10024		<i>mat A</i>	Portugal, Tapada de Mafra
10027		<i>mat a</i>	Portugal, Monchique
10028		<i>mat a</i>	Portugal, Monchique
10036		<i>mat a</i>	Italy, Turchino Est.
10037		<i>mat a</i>	Italy, Turchino Est.
10038		<i>mat a</i>	Italy, Turchino Est.
10040		<i>mat a</i>	Italy, Turchino Est.
10042		<i>mat a</i>	Italy, Turchino Est.
10051		<i>mat A</i>	Italy, Genoa
10054		<i>mat a</i>	Italy, Genoa
10056		<i>mat A</i>	Italy, Genoa
<i>N. discreta</i>			
9991	D221	<i>mat A</i>	Spain, Macanet de la Selva
9990	D220	<i>mat A</i>	Portugal, Monchique
9989	D218	<i>mat A</i>	Portugal, Monchique
10025		<i>mat a</i>	Portugal, Monchique
10010		<i>mat a</i>	Portugal, Boticas
10011		<i>mat a</i>	Portugal, Boticas
9986	D215	<i>mat a</i>	Portugal, Boticas
9987	D216	<i>mat A</i>	Portugal, Boticas
9988	D217	<i>mat a</i>	Portugal, Boticas
10012		<i>mat a</i>	Portugal, Boticas
10013		<i>mat A</i>	Portugal, Boticas
10014		<i>mat A</i>	Portugal, Boticas
9992	D224	<i>mat A</i>	Switzerland, Leuk
9993	D225	<i>mat A</i>	Switzerland, Leuk

^aD numbers refer to isolate numbers given by Dettman et al (2006) in the phylogenetic study of *N. discreta*.

Characterization of the genetic diversity among N. crassa and N. discreta strains.—Phylogenetic analyses of *N. crassa* and *N. discreta* have revealed genetically distinct clades within these species (Dettman et al 2003a, 2006). To assign European isolates to these clades, or to discover other clades within these biological species, sequence was obtained for three diagnostic polymorphic DNA regions (Dettman et al 2003a). Sequences of the three polymorphic regions (unlinked, noncoding loci that flank microsatellites [TMI, TML, and DMG]) were obtained with methods described by Dettman et al (2003a). Sequences were aligned manually, because of the presence of microsatellites and insertion/deletion gaps (indels) within these loci. Microsatellite sequences

were omitted from the analyses. Fourteen of 17 European *N. discreta* isolates were analyzed (excluding multiple isolates of the same mating type from the same plant), as were 22 *N. crassa* isolates from all sites where *N. crassa* was present, including multiple isolates of different mating type where available (TABLE I). The only two *N. crassa* isolates obtained from western North America (Montana; FGSC 8571 and W-864) (Jacobson et al 2004) also were included. The sequences have been deposited in GenBank under accession numbers DQ442288–DQ442377.

The sequences of the three loci were combined into a single dataset because previous use of the partition homogeneity test showed a lack of incongruence (Dettman

et al 2003a, 2006). Separate maximum parsimony trees were calculated for *N. discreta* and *N. crassa* with PAUP* (version 4.0b10, Swofford 2003). Analysis of European *N. discreta* isolates in relation to worldwide collections of *N. discreta sensu lato* has been reported by Dettman et al (2006). For comparative purposes the *N. crassa* dataset included sequences of the three loci from a subset of 37 of the *N. crassa* strains included in Dettman et al (2003a). No outgroups were included, because Dettman et al (2003a) clearly showed that *N. crassa* is a well supported phylogenetic species. Maximum parsimony bootstrapping for *N. crassa* was performed with heuristic searches (1000 replicates, simple stepwise addition, tree bisection-reconnection branch swapping, MAXTREES = 100).

RESULTS

The occurrence of Neurospora in wildfire sites.—The yellow to orange colonies of *Neurospora* conidiating on the surface of woody and herbaceous plants killed by fire were recognized easily (FIG. 1). *Neurospora* was found to be common at some sites, while being relatively rare at others sites (TABLE II, FIG. 2). The 14 sites surveyed extend over ca. 1650 km in a path leading generally northeast from southern Portugal (37°18'N, 8°35'W) to Switzerland (46°19'N, 7°38'E). The number of isolates collected totals 247 and includes the species *N. crassa*, *N. discreta*, *N. sitophila* and *N. tetrasperma*. At five of the 14 sites collections were from a single species of *Neurospora*, whereas collections at the other nine sites yielded multiple species. Most of these isolates (195) were single colonies collected from an individual plant. Multiple colonies (2–7) were sampled from 13 plants across five sites. Eleven of these plants yielded a single species, but two plants were colonized by two *Neurospora* species each. Although no systematic attempt was made to gauge the level of clonality or measure intraspecific genetic diversity among the isolates, multiple genotypes of the same species were found on five of the plants inhabited by a single species of *Neurospora* (see below).

Spore killer in N. sitophila.—Isolates identified as *N. sitophila* could be separated into two classes based on crosses with the species tester strains. One class produced 90–95% black ascospores, whereas the other produced 50% black ascospores with the remaining spores being hyaline, significantly smaller and unviable. When perithecia from these crosses were dissected microscopically, nearly every ascus showed a 4:4 black:hyaline ascospore pattern (FIG. 3). This pattern is the hallmark of Spore killer meiotic drive in *Neurospora* (Raju



FIG. 1. *Neurospora* growing and sporulating on scorched vegetation in Europe. A. Extensive colonization of an unidentified shrub at Turchino Est., Italy. B. Localized sporulation limited to the node of cane-like grass at Seros, Spain.

2002). Because the tester strains of *N. sitophila* used (FGSC 5940 *mat A* and 5941 *mat a*) are known to be sensitive to Spore killer, the killer component must be present in the European *N. sitophila* isolates.

A single spore killer element, Spore killer (*Sk-1*), has been described in *N. sitophila* (Raju 2002, Turner 2001). Research with *Sk-1* has shown that only killer × sensitive heterozygous crosses show killing; both homozygous crosses, killer × killer and sensitive × sensitive, show normal 8:0, black:hyaline, ascospores in each ascus. Therefore, to determine if the Spore killer in European isolates is *Sk-1* or a new element, each *N. sitophila* strain was crossed to *Sk-1* testers strains (FGSC 2216 *mat A* or 2217 *mat a*). All European isolates that showed 4:4 killing when crossed to sensitive produced 8:0 asci when crossed to *Sk-1*. Likewise all European isolates that produced 8:0 asci when crossed to sensitive showed 4:4 killing when crossed to *Sk-1*. All killer European *N. sitophila* isolates, therefore, are *Sk-1*; no new killer elements were apparent in these samples.

Fifty-four percent of the *N. sitophila* isolates collected (45 of 83) expressed the killer phenotype. The killer haplotype was present in six of the seven sites containing *N. sitophila*, and three of these contained both killer and sensitive haplotypes. However killer and sensitive haplotypes were not found together on any of the five plants from which multiple isolates of *N. sitophila* were recovered.

N. crassa.—A single maximum parsimony tree was produced from combined sequences of the TMI, TML, and DMG loci (FIG. 4). Included in the tree were representatives of the three major clades in

TABLE II. The distribution of species of *Neurospora* across sites surveyed in Europe in 2003–2004^a

Country	Site	Latitude	Longitude	<i>N. crassa</i>		<i>N. sitophila</i> ^b		<i>N. discreta</i>		<i>N. tetrasperma</i>
				<i>matA</i>	<i>mata</i>	<i>matA</i>	<i>mata</i>	<i>matA</i>	<i>mata</i>	<i>matA+a</i>
Portugal	Monchique	37°18'	8°35'W		2		12 (0:12)	5	1	
	Tapada de Mafra	38°58'	9°17'W	2	1	3 (3:0)				1
	Penedo Furado	39°38'	8°10'W	1						1
	Troviscal Sertã	39°52'	8°0'W	1						4
	Boticas	41°42'	7°41'W					3	5	1
Spain	Sapiãos	41°43'	7°37'W							10
	Sevilla	37°24'	5°59'W	3	23					
	Seros	41°23'	0°19'E	20	22					
	Macanet de la Selva	41°46'	2°43'E		1	2 (2:0)	5 (4:1)	1		
	Platja d'Aro	41°50'	3°4'E	6	4	2 (2:0)				
France	Vidauban	43°24'	6°28'E							15
Italy	Genova	44°26'	8°45'E	4	20	22 (17:5)	8 (8:0)			
	Turchino Est.	44°27'	8°44'E		5	14 (0:14)	9 (3:6)			
Switzerland	Leuk	46°19'	7°38'E			6 (6:0)		2		
Totals				37	78	49	34	11	6	
% of total (247)					115 46%	83 34%		17 7%		32 13%

^aAll isolates are totaled here, including those collected from the same plant. No systematic attempt was made to identify clones which may have been repeatedly sampled. Characterization of a small number of genetic markers, for a limited number of isolates, was conducted for phylogenetic clade identification and tree construction (see TABLE I, FIG. 4 and text).

^bThe ratio of *Sk-I* killer to sensitive isolates is in parentheses.

N. crassa, NcA, NcB and NcC (Dettman et al 2003a). Sequence was obtained for all three loci from 22 European isolates of *N. crassa*. All these isolates fell into the single, previously described clade NcB (TABLE I, FIG. 4).

The sequence of the TMI locus subsequently was obtained from the remaining 93 isolates of European *N. crassa* to associate each with the appropriate phylogenetic clade. TMI was chosen as a diagnostic locus because its sequence is clearly distinct between the NcB clade versus clades NcA and NcC. All 93 isolates fell within the NcB clade (data not shown). Of these 83 had TMI sequences that were essentially identical, including the number of microsatellite repeats. Ten isolates, all from Seville, Spain, were exceptional in having a single nucleotide polymorphism at base 119 in the microsatellite flanking sequence and a microsatellite with 5 rather than 12 repeats.

Although it was beyond the scope of this study to assess clonality of strains from the same plant, we did investigate the genotypes of multiple *N. crassa* isolates

collected from seven individual plants. When two polymorphic markers (*mat* and TMI) were combined, five plants from Seville, Spain, revealed more than one genetically distinct individual of *N. crassa* per plant. In contrast the multiple isolates of *N. crassa* from the two other plants (from Seros, Spain, and Genoa, Italy) were monomorphic at both markers. This preliminary study indicated that more than one genetic individual could be present in very close spatial scales, as was reported by Powell et al (2004).

Phylogenetic species 4B within the N. discreta complex.—The European isolates of *N. discreta sensu lato*, as defined by biological species recognition, all were identified as belonging to phylogenetic species (PS) 4B (tree not shown, refer to Dettman et al 2006 FIG. 2 for relationships among phylogenetic species within the *N. discreta* complex). Of the six isolates sequenced here that were not analyzed by Dettman et al (2006), each had sequence identical to at least one isolate examined




FIG. 2. Distribution of *Neurospora* biological species collected in Europe.

by Dettman et al (2006). Therefore no additional genetic diversity was found within PS 4B or the European population, and PS 4B is the only species of the *N. discreta* complex found in Europe to date.

DISCUSSION

Reports of the occurrence of *Neurospora* in Europe have been published sporadically over the past 160 y, beginning with its earliest description from France in 1843 (see Perkins 1991). Most of these descriptions have concentrated on *Neurospora* contamination of bakeries and their products; the most recent was Yassin and Wheals (1992). Not long after formal description of the genus by Shear and Dodge (1927), however, Ramsbottom and Stephens (1935) mentioned that *Neurospora* was found on other natural substrates, most notably burnt trees and gorse in Britain. Other anecdotal observations have suggested that *Neurospora* is not uncommon in Europe (e.g. D. Zickler, University Paris South, personal communication with D.D. Perkins, Stanford University). However, to our knowledge, this is the first study that systematically sampled *Neurospora* from natural habitats in Europe.

All five classically described, conidiating, biological species of *Neurospora* now have now been identified in Europe: *N. crassa*, *N. discreta*, *N. intermedia*, *N. sitophila* and *N. tetrasperma*. This is the first study to report *N. discreta*, whereas we did not find *N. intermedia*, which was reported by Ramsbottom and Stephens (1935). Recent work has further divided *Neurospora* into phylogenetic species and clades (Dettman et al 2003a, 2006). Of the eight phylogenetic species within the *N. discreta* complex only one (PS 4B) was identified among the European isolates




FIG. 3. Asci from crosses of European *N. sitophila* Spore killer strains. A. A cross heterozygous for *Sk-1* (killer \times sensitive). Asci contain four normal size, maturing, viable *Sk-1* ascospores and four hyaline, aborted sensitive ascospores. B. A homozygous (sensitive \times sensitive) cross for *Sk-1*. Asci contain eight normal size, maturing, viable ascospores. Homozygous *Sk-1*killer \times killer crosses also show asci containing eight viable ascospores (Photomicrographs courtesy of N.B. Raju, Stanford University).

collected here. Two newly described phylogenetic species outside the *N. discreta* complex also were found to be distinct biological species (Dettman et al 2003b); neither of these species were found among the European isolates. Of the three distinct clades within *N. crassa* (NcA, NcB and NcC), all new European *N. crassa* isolates fell into NcB. Based on these finer scale measures of genetic divergence among members of *Neurospora*, similarities and differences were assessed between the newly sampled populations from Europe and populations from other continents, including both temperate and tropical/subtropical climates.

The similarity of *Neurospora* between Europe and southeastern, subtropical areas of the United States is also reflected in the overall species diversity and distributions. The complement of species and their frequency of collection are similar in Europe and southeastern United States (FIG. 5). This distribution is in stark contrast to populations of *Neurospora* in western North America, which are composed pre-




FIG. 4. The relationships among the three phylogenetic clades within *N. crassa*. Maximum parsimony, unrooted phylogram produced from the sequences of three combined loci. Numbers next to bold branches separating NcA, NcB and NcC clades indicate bootstrap support (1000 replicates). Taxon labels indicate strain number and geographic source; European strains from this study are shown within the box, and western North American strain from Jacobson et al (2004) are shown in the shaded box. All strains labeled with D numbers were sequenced as part of Dettman et al (2003a). One locus (TMI) was sequenced from an additional 93 European *N. crassa* isolates because this locus is diagnostic for *N. crassa* clade. All 93 isolates were placed definitively into NcB based on TMI sequence (data not shown).

dominantly of a single species in the *N. discreta* complex (PS 4B), with only rare occurrences of *N. sitophila* and *N. crassa*. *Neurospora* has been found at 64°N latitude in Alaska and as far as 45°N in Europe. Future collecting expeditions are planned to target even higher latitudes in Europe in the hope of learning more about the distribution of *Neurospora* species.

The absence of *N. intermedia* in our European

collection was unexpected given reports in the literature (Ramsbottom and Stephens 1935, Yassin and Wheals 1992). For example Yassin and Wheals (1992) reported nine of 345 isolates (<3%) as *N. intermedia*, eight of which were from nonbakery sources, including imported Indonesian onjom. Moreover *N. intermedia* is by far the most common species collected world wide, particularly at latitudes >30°N and S (China, Japan, Australia and New




FIG. 5. Frequency of *Neurospora* biological species by region. Data for western North America taken from Jacobson et al (2004), data for southeastern United States taken from Turner (2001).

Zealand) (Turner et al 2001). Given that there are likely to be sources of *N. intermedia* in Europe, the lack of *N. intermedia* in our collections from temperate northern latitudes in Europe and western North America is intriguing but its significance cannot be assessed currently.

The physical appearance in nature of *N. crassa* and *N. discreta* from Europe and the southeastern US is remarkably similar and unlike that of *N. discreta* from western North America. However phylogenetic analysis of DNA sequences indicated that the European isolates of the two species were highly similar to those from temperate western North America and dissimilar to those found in the southeastern United States.

N. crassa clades NcA, NcB and NcC are genetically distinct from one another but do not meet the strict criteria that would make them separate phylogenetic species (Dettman et al 2003a). These clades have distinct geographical distributions. NcA was widespread across the Caribbean basin and Africa. NcC was limited to the state of Tamil Nadu in India, and the rare isolates of NcB were limited to equatorial Africa and southern India. The addition of all the European and western North American isolates of *N. crassa* to clade NcB significantly changes the biogeography of the species. Now NcB also appears geographically widespread, similar to NcA, although its prevalence outside of Europe remains in question.

Distributions of NcA and NcB in the western hemisphere and Europe are nonoverlapping, but the clades do coexist in equatorial Africa. NcA and NcC, whose ranges overlap in southern India, have developed reproductive isolation phenotypes, which

correlate with the genetic distance (Dettman et al 2003b; E. Turner, University of California at Berkeley, unpublished). No attempt was made in this or previous studies to characterize the reproductive relationships between members of the NcA and NcB clades; biological species recognition was limited to crossing European isolates to the species tester strains. We therefore do not know whether NcA and NcB show reproductive isolation anywhere in their range.

The *N. discreta* complex from Europe and North America also shows a combination of widespread and more narrowly distributed species. European *N. discreta* isolates, which represent only 7% of all collected European isolates, are placed phylogenetically within the predominant species in western North America, the widespread PS 4B. PS 4B however is phylogenetically distant from the two other *N. discreta* species in North America (i.e., *N. discreta sensu stricto* [Texas] and PS 7 [Florida, Mexico and Guatemala]).

Striking differences were seen in the growth habit and the morphology of colonies of *Neurospora* on natural substrates between the two temperate continents. As mentioned, *Neurospora* in Europe was commonly seen apparently growing on the surface of charred bark (FIG. 1A). In contrast extensive colonies of *Neurospora* were seen in western North America but always under the bark of woody plants. Only rarely, and after prolonged incubation periods, did the fungus erupt through the bark (Jacobson et al 2004, FIG. 1). The extensive amount of sporulation on the surface of burned bark, as seen in Europe, was never observed in western North America. Moreover recognizable colonies of *Neurospora* were not observed under the bark in Europe.

Neurospora in Europe grew on both charred woody and herbaceous plants, such as the grass in Spain (FIG. 1B). In contrast *Neurospora* was never observed in western North America on herbaceous plants. Although the latitude, climate, geography and vegetation are similar between Europe and temperate western North America, the growth habit and substrate of *Neurospora* in Europe are similar to those in tropical and subtropical areas, including Florida and Texas in the southern United States (Powell et al 2003, Turner and Perkins 1988, Turner et al 2001).

The proportion of *N. sitophila* Spore killer strains reported here for Europe is much higher (45 of 83, 54%) than worldwide (77 of 469, 16%) (FIG. 5) and might provide an opportunity to study the dynamics of Spore killers in nature and the effect of meiotic drive on populations. Existing data have been insufficient to determine the potential of Spore killer to become fixed in any population (Turner 2001), which makes the spatial and temporal dynamics of killer and

sensitive haplotypes in European populations of great interest (Burt and Trivers 2006). Whether a stable equilibrium is maintained can be tested only where killer and sensitive coexist in the same populations, but *Sk-I* killer and sensitive haplotypes coexist from only 10 out of 92 (11%) sites where Spore killer has been found outside of Europe: one in Hawaii, one in Vanuatu and eight in Tahiti. In Europe, as mentioned, killer and sensitive coexist in three of six sites with *Sk-I* frequency of 13–83%. European Spore killer isolates also were reported by Yassin and Wheals (1992) who found that all nine of their bakery *N. sitophila* isolates were *Sk-I*. Re-sampling of European *N. sitophila* populations over time might provide the data needed to understand Spore killer and meiotic drive in nature.

Together with the recent discovery of *Neurospora* in western North America (Jacobson et al 2004), documentation presented here of its occurrence in Europe firmly establishes it as a common inhabitant of temperate climates, perhaps worldwide. The broad distribution of *N. discreta*, particularly its longitudinal component, place it, along with *N. crassa*, among the handful of species that have the attributes to serve as evolutionary and ecological model organisms. There will be no lack of ecological questions because of the large gaps in our knowledge of the basic ecology of *Neurospora* and fire adapted fungi in general. The differences in regional diversity, reported here, can form the basis of testable hypotheses. Questions of phylogeography and adaptations specifically can be addressed with *Neurospora*: Where did these species of *Neurospora* originate and how did they arrive at their modern distributions? What role have human activities played in the current distribution of *Neurospora* lineages? Have populations of *Neurospora* changed genetically to adapt to local conditions, such as ambient temperature or photoperiod (Tan et al 2004)? How organisms spread and adapt to new environments are fundamental ecosystem properties, yet they are little understood. We hope that understanding of these fundamental features will come from studies of *N. discreta* that blend ecology and evolutionary biology with genetics and genomics.

ACKNOWLEDGMENTS AND CONTRIBUTIONS OF AUTHORS

D.J. conceived and coordinated the project with the assistance of J.T., M.M., T.R., A.V., and L.M.C. D.J., C.B. and S.S. collected and cultured most strains. M.D., I.M., A.U., P.C. and A.V. collected and cultured Portugal strains, identifying some. L.M.C., L.N.S. and M.O. collected in Seville and provided needed logistical help in locating other collection sites in Spain. D.J. identified or confirmed identification of all strains. J.D., R.A. and D.J. sequenced and performed phylogenetic analyses. D.J. analyzed the data

and with J.T. wrote the paper. The work was supported by a grant from the US National Science Foundation to J.T. (DEB-0316710); by grants from the Deutsche Forschungsgemeinschaft and the Meyer-Struckman Stiftung to T.R.; by grants from Fundação para a Ciência e a Tecnologia to A.V.; and by grants from the Ministerio de Educación y Ciencia, Spain, (INIA RM2004-00007) and Junta de Andalucía (CVI 0119) to L.M.C. D.J. also is supported in part by US National Science Foundation grant MCB-0417282, awarded to David D. Perkins, Stanford University. We thank David Perkins for allowing part of this work to be completed in his laboratory and N.B. Raju for the photomicrographs used in FIG. 3.

LITERATURE CITED

- Burt A, Trivers R. 2006. Genes in conflict: the biology of selfish genetic elements. Cambridge, Massachusetts: Belknap Press of Harvard University Press. 602 p.
- Cai L, Heewon R, Hyde KD. 2006. Phylogenetic investigations of *Sordariaceae* based on multiple gene sequences and morphology. *Mycol Res* 110:137–150.
- Davis RH. 2000. *Neurospora*: contributions of a model organism. New York: Oxford University Press. 333 p.
- Dettman JR, Jacobson DJ, Taylor JW. 2003a. A multilocus genealogical approach to phylogenetic species recognition in the model eukaryote *Neurospora*. *Evolution* 57:2703–2720.
- , ———, ———. 2006. Multilocus sequence data reveal extensive phylogenetic species diversity within the *Neurospora discreta* complex. *Mycologia* 98:437–447.
- , ———, Turner E, Pringle A, Taylor JW. 2003b. Reproductive isolation and phylogenetic divergence in *Neurospora*: comparing methods of species recognition in a model eukaryote. *Evolution* 57:2721–2741.
- García D, Stchigel AM, Cano J, Guarro J, Hawksworth DL. 2004. A synopsis and recircumscription of *Neurospora* (syn. *Gelasinospora*) based on ultrastructural and 28S rDNA sequence data. *Mycol Res* 108:1119–1142.
- Jacobson DJ, Powell AJ, Dettman JR, Saenz GS, Barton MM, Hiltz MD, Dvorachek WH, Glass NL, Taylor JW, Natvig DO. 2004. *Neurospora* in temperate forests of western North America. *Mycologia* 96:66–74.
- Legan JD. 1993. Mould spoilage of bread: the problem and some solutions. *Int Biodeterior Biodegrad* 32:33–53.
- Perkins DD, Turner BC. 1988. *Neurospora* from natural populations: toward the population biology of a haploid eukaryote. *Exp Mycol* 12:91–131.
- . 1991. The first published scientific study of *Neurospora*, including a description of photoinduction of carotenoids. *Fung Genet Newsl* 38:64–65.
- Powell AJ, Jacobson DJ, Natvig DO. 2003. Variation among natural isolates of *Neurospora* on small spatial scales. *Mycologia* 95:809–819.
- Raju NB. 2002. Spore killers: meiotic drive elements that distort genetic ratios. In: Osiewacz HD, ed. *Molecular biology of fungal development*. New York: Marcel Dekker Inc. p 275–296.
- Ramsbottom J, Stephens FL. 1935. *Neurospora* in Britain. *Trans Brit Mycol Soc* 19:215–220.

- Shear CL, Dodge BO. 1927. Life histories of and heterothallism of the red bread-mold fungi of the *Monilia sitophila* group. *J Ag Res* 34:1019–1042.
- Swofford DL. 2003. PAUP*: phylogenetic analysis using parsimony (*and other methods). Version 4.0.b10. Sunderland, Massachusetts: Sinauer Associates.
- Tan Y, Merrow M, Roenneberg T. 2004. Photoperiodism in *Neurospora crassa*. *J Biol Rhythm* 19:135–43.
- Turner BC. 2001. Geographic distribution of *Neurospora* Spore killer strains and strains resistant to killing. *Fung Genet Biol* 32:93–104.
- , Perkins DD, Fairfield A. 2001. *Neurospora* from natural populations: a global study. *Fung Genet Biol* 32:67–92.
- Yassin S, Wheals A. 1992. *Neurospora* species in bakeries. *J Appl Bacteriol* 72:377–380.